Asian Journal of Multidimensional Research ISSN: 2278-4853 Vol. 12, Issue 6, June 2023 SJIF 2022 = 8.179 A peer reviewed journal

METHODS OF DIAGNOSIS AND TREATMENT OF ANEMIA IN HEMODIALYSIS PATIENTS

Makhmonov Lutfullo Saydullaevich*; Turdiev Qilich Toshkobilovich**; Elmirzaev Akhmad Boliboyevich***

> *Doctor, Head of Department of Hematology, SOMC, UZBEKISTAN

> > **PhD,

Head of Department of Hematology, Hemodialysis SOMMC, UZBEKISTAN

***Doctor-dep, Head of Department of Hematology, Hemodialysis SOMMC, UZBEKISTAN DOI: 10.5958/2278-4853.2023.00079.4

ABSTRACT

Anemia is an independent risk factor for the development of cardiovascular, infectious and septic complications in dialysis patients. Timely and complete correction of anemia to the recommended target indicators significantly reduces morbidity and mortality, as well as improves the quality of life and medical and social rehabilitation of this category of patients. In the article, a retrospective analysis of the effectiveness of the use of recormon in the correction of anemia in 389 patients with hemodialysis was performed over 5 years.

KEYWORDS: Anemia, Hemodialysis, Correction of Anemia, Erythropoiesis, Sexual Functions, Social Rehabilitation.

REFERENCES:

- 1. Ermolenko V.M., Ivashchenko M.A. Uremia and erythropoietin; 2010.
- **2.** Shilo V.Yu., Gorin A.A., Denisov A.Yu. Quality of life of patients on hemodialysis: relationship with the achievement of the target level of hemoglobin. Abstracts of the IV Conference of the RDO. Nephrology and Dialysis 2015;7(3):303.
- Fishbein S., Paganini E.P. Hematological anomalies. In: Guide to dialysis. Ed. D. Daugirdas, P. Blake, T. Ing. 3rd ed./Trans. from English. ed. A.Yu. Denisov and V.Yu. Awl. Tver; Triada, 2013. p. 528–46.
- **4.** Tareeva I.E., Nikolaev A.Yu., Klepikov P.V. Multicenter trial of Recormon in an international controlled study. Terarch 1992;64(8):59–62.
- **5.** Krisper P., Quehenberger F., Schneditz D. et al. Prediction of time-averaged concentration of haemoglobin in haemodialysis patients. Nephrol Dial Transplant 2013;18(10):2082–7.

- 6. NKF-DOQI Clinical Practice Guidelines for the treatment of anemia of chronic renal failure. National Kidney Foundation-Dialysis Outcomes Quality Initiative. Am J Kidney Dis 2007;(4 Suppl 3):P. 192–240.
- **7.** NKF-KDOQI Clinical Practice Guidelines for anemia of chronic kidney disease: update 2000. New-York, 2011.
- **8.** European best practice guidelines for the management of anemia in patients with chronic renal failure. Working Party for European Best Practice Guidelines for the Management of Anaemia in Patients with Chronic Renal Failure. Nephrol Dial Transplant 2009;14 (Suppl 5):1–50.
- **9.** Rumyantsev A.G., Morshchakova E.F., Pavlov A.D. Anemia in chronic renal failure. In: Erythropoietin in the diagnosis, prevention and treatment of anemia. M.; 2013. p. 211–55.
- **10.** Storozhakov G.I., Gendlin G.E., Tomilina N.A. etc. Defeat of the cardiovascular system in chronic renal failure. Ross Medical Journal 2015;(2):4–8.
- **11.** European Best Practice Guidelines for Hemodialysis (Part 1). Section IV: Dialysis fluid purity. Nephrol Dial Transplant 2002;17 (Suppl 7):45–62.
- **12.** Rybakova O.B., Denisov A.Yu., Shilo V.Yu. Correction of anemia in patients with terminal chronic renal failure: comparative effect of hemodialysis and on-line hemodiafiltration. Nephrology and Dialysis 2012;(4):270–4.
- **13.** Shilo V.Yu., Khasabov N.N. Anemia in patients on dialysis: iron deficiency, methods for its diagnosis and correction. Anemia 2014;(1):19–29.